

RESEARCH ARTICLE

WILEY

Cannabinoid receptor activation potential of the next generation, generic ban evading OXIZID synthetic cannabinoid receptor agonists

Marie H. Deventer¹  | Katleen Van Uytfanghe¹  | Inge M. J. Vinckier²  |
Fabiano Reniero³  | Claude Guillou³  | Christophe P. Stove¹ 

¹Laboratory of Toxicology, Department of Bioanalysis, Faculty of Pharmaceutical Sciences, Ghent University, Ghent, Belgium

²Laboratory of Customs and Excises, Vilvoorde, Belgium

³European Commission, Joint Research Centre, Directorate F-Health, Consumers and Reference Materials, Ispra, Italy

Correspondence

Christophe P. Stove, Laboratory of Toxicology, Department of Bioanalysis, Faculty of Pharmaceutical Sciences, Ghent University, Ottergemsesteenweg 460, 9000 Ghent, Belgium.

Email: christophe.stove@ugent.be

Funding information

Research Foundation-Flanders, Grant/Award Number: 1S54521N

Abstract

In recent years, several nations have implemented various measures to control the surge of new synthetic cannabinoid receptor agonists (SCRAs) entering the recreational drug market. In July 2021, China put into effect a new generic legislation, banning SCRAs containing one of seven general core scaffolds. However, this has driven manufacturers towards the synthesis of SCRAs with alternative core structures, exemplified by the recent emergence of “OXIZID SCRAs.” Here, using *in vitro* β -arrestin2 recruitment assays, we report on the CB₁ and CB₂ potency and efficacy of five members of this new class of SCRAs: BZO-HEXOXIZID, BZO-POXIZID, 5-fluoro BZO-POXIZID, BZO-4en-POXIZID, and BZO-CHMOXIZID. All compounds behaved as full agonists at CB₁ and partial agonists at CB₂. Potencies ranged from 84.6 to 721 nM at CB₁ and 2.21 to 25.9 nM at CB₂. Shortening the n-hexyl tail to a pentyl tail enhanced activity at both receptors. Fluorination of this pentyl analog did not yield a higher receptor activation potential, whereas an unsaturated tail resulted in decreased potency and efficacy at CB₁. The cyclohexyl methyl analog BZO-CHMOXIZID was the most potent compound at both receptors, with EC₅₀ values of 84.6 and 2.21 nM at CB₁ and CB₂, respectively. Evaluation of the activity of a seized powder containing BZO-4en-POXIZID suggested a high purity, in line with high-performance liquid chromatography coupled to diode-array detection (HPLC-DAD), gas chromatography coupled to mass spectrometry (GC-MS), liquid chromatography coupled to time-of-flight mass spectrometry (LC-QTOF-MS), and Fourier transform infrared spectroscopy (FTIR) and nuclear magnetic resonance (NMR) analysis. Furthermore, all tested compounds showed a preference for CB₂, except for BZO-POXIZID. Overall, these findings inform public health officials, law enforcement agencies, and clinicians on these newly emerging SCRAs.

KEYWORDS

bioassay, CB₁ cannabinoid receptor, new psychoactive substances, OXIZID, synthetic cannabinoid receptor agonists

1 | INTRODUCTION

Synthetic cannabinoid receptor agonists (SCRAs) remain one of the most identified classes of new psychoactive substances (NPS) worldwide, and their number continues to increase.¹ Eleven new SCRAs were reported in Europe for the first time in 2020, adding up to a total of 209 compounds being detected since 2008.² Although a decrease in the number of newly detected SCRAs has been noticed during the last years,¹ monitoring SCRA use remains important in specific settings, for instance among homeless people and in prisons (in Europe), the latter usually to circumvent mandatory drug tests.^{3,4} SCRAs exert their main sought-after psychoactive effects at the CB₁ cannabinoid receptor, thereby mimicking the effects of the phytocannabinoid Δ^9 -tetrahydrocannabinol (THC), the primary psychoactive substance of cannabis. Their major threat lies in their often much higher potency and efficacy compared with THC,⁵⁻⁷ the variability in the composition of the marketed products, and their easy accessibility via the Internet.⁸ Additionally, there has been an increase in reports of cannabis adulterated with potent SCRAs, resulting in users being unaware of the potential harms they could be exposed to.² SCRAs have been

associated with psychosis, agitation, hallucinations, seizures, respiratory failure, cardiovascular effects, coma, and even death.⁹⁻¹³

SCRAs and NPS in general are hard to put under legislative control, as unknown substances appear on the illicit drug market at a rapid pace. The stabilizing number of new NPS detected during recent years suggests the impact of regulatory steps taken by several nations, such as the introduction of generic legislations. These allow a nation to ban a larger group of substances encompassing certain core structures.^{1,17} However, the already complex recreational drug market remains a “game of cat and mouse,” as clandestine labs manage to find “legal loopholes” by synthesizing structurally diverse compounds not covered by the current control measures. For instance, in May 2021, the Office of China National Narcotics Control Commission announced that as of July 1, 2021, a generic legislation would be in place to control synthetic cannabinoids, similar to the scheduling of fentanyl-related substances in 2019.^{18,19} As opposed to the former individual listing of substances, which required the cumbersome identification and specification of each individual compound to be controlled, this measure allowed for a nationwide ban of compounds structurally related to seven general scaffolds (see Figure 1).¹⁴ This




FIGURE 1 Overview of the seven general SCRA scaffolds covered by the generic control measure in China, in effect as of July 1, 2021. The figure is based on the official announcement document, released by the Office of China National Narcotics Control Commission on May 12, 2021¹⁴

TABLE 1 Comparison of different nomenclature for the discussed substances

| Initial naming | Synonyms | IUPAC naming | New systematic naming |
|-------------------|-----------------------------------|---|-------------------------------|
| MDA-19 | MDA19 MDA 19 | (Z)-N-(1-hexyl-2-oxoindolin-3-ylidene)benzohydrazide | BZO-HEXOXIZID |
| Pentyl-MDA-19 | 5C-MDA-19 MDA-19 pentyl analog | (Z)-N-(1-pentyl-2-oxoindolin-3-ylidene)benzohydrazide | BZO-POXIZID |
| 5F-MDA-19 | MDA-19 5-fluoropentyl analog | (Z)-N-(1-(5-fluoropentyl)-2-oxoindolin-3-ylidene)benzohydrazide | 5F-BZO-POXIZID |
| 4en-pentyl MDA-19 | | (Z)-N'-(2-oxo-1-(pent-4-en-1-yl)indolin-3-ylidene)benzohydrazide | BZO-4en-POXIZID ¹⁵ |
| CHM-MDA-19 | Cyclohexylmethyl MDA-19 | (Z)-N-(1-(cyclohexylmethyl)-2-oxoindolin-3-ylidene)benzohydrazide | BZO-CHMOXIZID |

Note: The table is based on the Public Health Alert Report, prepared by NPS Discovery (CFSRE) and Cayman Chemical, and released on August 31, 2021.¹⁶

may have driven manufacturers towards the synthesis of compounds with new core structures, not covered by this legislation.¹⁵ The first series of such compounds was recently reported by Liu et al., who identified and characterized AD-18, 5F-MDA-19, and pentyl-MDA-19 in a seizure of powders and e-liquids.²⁰ As the term MDA (stemming from M.D. Anderson Cancer Center, the Institute where these compounds were first synthesized—see below) may be confused with the abbreviation of methylenedioxyamphetamine, a new, more “SCRA-friendly” OXIZID nomenclature, based on the chemical structure and IUPAC name, was developed by Cayman Chemical and the NPS Discovery program at the US-based Center for Forensic Science Research & Education (CFSRE) (see Table 1).¹⁵ The term OXIZID refers to the OXoIndoline core attached to the aZIDe linker and will be used throughout this article. As these OXIZIDs introduce a new class of non-scheduled compounds, it is anticipated that the number of OXIZID SCRAs may further increase in the near future.

Unlike CB₁, which is predominantly present in the central nervous system, CB₂ is primarily located in cells of the immune system and is involved in inflammatory processes.^{21,22} As CB₂-selective agonists are believed to be devoid of the undesirable side effects correlated with CB₁ activation and, importantly, are not believed to be psychoactive, they are considered to be potentially interesting therapeutic tools.^{23–26} It is in this context that the CB₂ agonist MDA-19 (BZO-HEXOXIZID) was developed in 2008 and was later pharmacologically characterized at the University of Texas M.D. Anderson Cancer Center.^{27,28} It served as a lead compound in the development of therapeutics for the treatment of neuropathic pain, an often difficult-to-treat condition caused by trauma or disease of the somatosensory nervous system^{29,30} as a result of, for example, diabetic neuropathy, multiple sclerosis, trigeminal neuralgia, and postherpetic neuralgia.^{27,31}

The scarcely available literature also mentions *in vitro* antiproliferative potential of BZO-HEXOXIZID, as studied using melanoma, osteosarcoma, and hepatocellular carcinoma cell lines.^{32–34}

In October 2016, BZO-HEXOXIZID was notified for the first time in Spain and reported to the European Monitoring Centre for Drugs and Drug Addiction (EMCDDA).³⁵ In October 2021, the CFSRE reported on the seizure of BZO-HEXOXIZID and two of its analogs, the truncated BZO-POXIZID (pentyl-MDA-19) and its fluorinated counterpart 5F-BZO-POXIZID (5F-MDA-19) in the United States.^{36–38} Shortly after, the first appearance on the recreational drug market in China of BZO-POXIZID and 5F-BZO-POXIZID was reported by Liu et al.²⁰ Only 1 month later, the CFSRE reported on the identification in plant-like material of yet another analog, BZO-CHMOXIZID (CHM-MDA-19, with a cyclohexyl methyl moiety instead of the hexyl tail of MDA-19).³⁹ Most recently, in December 2021, the Hungarian government announced that BZO-POXIZID, BZO-CHMOXIZID, and another analog, BZO-4en-POXIZID, carrying an unsaturated tail seen in other commonly identified SCRAs such as MDMB-4en-PINACA and ADB-4en-PINACA,^{40–42} were to be added to the list of controlled NPS (Decree No 55/2014).^{43,44} Relatively little is known about the pharmacology and structure–activity relationship of these compounds.

The fact that BZO-HEXOXIZID and its related compounds have repeatedly been identified in seized materials suggests that these compounds might still possess the potential to activate CB₁ and may be used for their psychoactive properties. In this study, we therefore assessed the intrinsic receptor activation potential of a panel of five OXIZID SCRAs (BZO-HEXOXIZID, BZO-POXIZID, 5F-BZO-POXIZID, BZO-4en-POXIZID, and BZO-CHMOXIZID) at both CB₁ and CB₂ by means of activity-based bioassays, monitoring β -arrestin2 (β arr2) recruitment to the activated receptor (structures of the compounds can be found in Figure 2). Furthermore, these bioassays were also used to evaluate the CB_{1/2} receptor activation potential of a powder that was intercepted by the Belgian Customs in November 2021 and that was confirmed to contain BZO-4en-POXIZID.




FIGURE 2 Chemical structures of the OXIZIDs evaluated in this report, together with the reference compound CP55,940 and the prototypic SCRA JWH-018. Structures were made with the ChemDraw 19 professional software

2 | MATERIALS AND METHODS

2.1 | Materials and reagents

Dulbecco's modified Eagle's medium (DMEM) (GlutaMAX™), Opti-MEM I Reduced Serum, penicillin, streptomycin, and amphotericin B were procured from Thermo Fisher Scientific (Waltham, MA, USA). Fetal bovine serum (FBS) and poly-D-lysine were supplied by Sigma-Aldrich (Darmstadt, Germany). Methanol was purchased from Chem-Lab NV (Zedelgem, Belgium). BZO-HEXOXIZID, BZO-POXIZID, 5F-BZO-POXIZID, BZO-4en-POXIZID, and BZO-CHMOXIZID were kindly provided by Cayman Chemical (Ann Arbor, MI, USA). The Nano-Glo® Live Cell reagent and the Nano-Glo® LCS Dilution buffer were obtained from Promega (Madison, WI, USA). A powder sample containing BZO-4en-POXIZID was seized in November 2021 by the Laboratory of the Belgian Customs and Excise services (Vilvoorde, Belgium). All reagents used for the analytical characterization were at least of high-performance liquid chromatography (HPLC) grade. LC-MS grade methanol and formic acid were purchased from Chem-Lab NV. Acetonitrile was procured from Biosolve (Valkenswaard, The Netherlands). Ammonium formate, ortho-phosphoric acid (85%) and potassium dihydrogen phosphate were purchased from Sigma-Aldrich (Diegem, Belgium). For NMR analysis, all reagents were obtained from Sigma-Aldrich. For FTIR analysis, no solvents or reagents were used.

2.2 | In vitro CB₁ and CB₂ β-arrestin2 recruitment assays

To determine activity at CB₁ and CB₂, previously described live cell-based bioassays monitoring agonist-induced recruitment of the intracellular βarr2 protein to the activated receptor were used. The concept is based on the NanoLuc Binary Technology (NanoBiT®, Promega), monitoring the interaction between one inactive subunit of a nanoluciferase fused to the receptor, and the other subunit, fused to βarr2. Receptor activation by a ligand results in the recruitment of βarr2, bringing the two subunits in close proximity, resulting in functional complementation of the enzyme. In the presence of the furimazine substrate, measurable bioluminescence is generated. The development of the system and the establishment of the stable cell lines used for these assays have been reported before.^{45–47}

Human embryonic kidney (HEK) 293T stably expressing the CB₁-βarr2 or CB₂-βarr2 system were routinely maintained at 37°C, 5% CO₂ under humidified atmosphere in DMEM (GlutaMAX™), supplemented with 10% heat-inactivated FBS, 100 IU/ml of penicillin, 100 µg/ml of streptomycin, and 0.25 µg/ml of amphotericin B.

Stock solutions were made in MeOH. Working solutions were prepared by serial dilution in Opti-MEM containing 50% MeOH and were used within 24 h upon preparation.

On the day prior to the assay, cells were seeded in poly-D-lysine coated white opaque-walled 96-well plates at approximately 50,000 cells per well and left to incubate overnight. Next, to remove residual traces of serum that could potentially interfere with protein

interactions during the assay, cells were rinsed twice with 150 µl of Opti-MEM I reduced serum, and 100 µl of this medium was added to each well. The Nano-Glo® Live Cell reagent, a non-lytic cell reagent containing the furimazine substrate, was diluted 20-fold in LCS buffer, and 25 µl of this mix was added to each well. The plate was then placed in the TriStar² LB 942 Multimode Microplate Reader (Berthold Technologies GmbH & Co., Germany), and luminescence was monitored for 10–15 min during an initial equilibration phase, which will later be used to correct for *inter-well* variability. Upon stabilization of the signal, 10 µl of a 13.5× concentrated stock solution was added, and luminescence was measured during 2 h. A concentration range of the reference compound CP55,940 and appropriate solvent controls for the analyzed compounds were included on each plate. CP55,940 was selected as a reference because it was previously used for the characterization of MDA-19 and structural analogs.^{27,28} To allow for a better comparison with earlier work, JWH-018 was also taken along, as it has often served as a reference compound in the used bioassays.^{48–50} All test concentrations were run in duplicate in minimally three independent experiments.

2.3 | Data analysis and statistical analyses

Raw data were processed using Microsoft Excel 2019, followed by curve fitting and statistical analysis using the GraphPad Prism software (Version 9.3.0) (San Diego, CA, USA). Firstly, to correct for inter-well variability, a baseline correction was performed on the absolute luminescence values, using data generated during the equilibration period. Then, for each compound, the mean area under the curve (AUC) was calculated. A blank correction was performed by subtracting AUC values of the solvent controls. Results represent the AUC ± standard error of mean (SEM) and were obtained by normalizing to the E_{max} of the reference compound CP55,940, arbitrarily set at 100%. Data points were consistently excluded for the highest concentration in case of a signal reduction of 20% or more compared to the next dilution, as this could potentially be a sign of cell toxicity or solubility issues at higher concentrations. Potency and efficacy were assessed by calculating pharmacological parameters EC₅₀ and E_{max} by curve fitting the obtained concentration–response curves via nonlinear regression (three-parameter logistic fit). Outliers were detected using the Grubbs test and omitted from the dataset if applicable (*p* value < 0.05; applicable for 1 out of 958 data points).

Receptor selectivity was evaluated and quantitated using an intrinsic relative activity-based method, commonly employed to calculate pathway bias.^{49,51–53} For each test compound, the intrinsic relative activity (RA_i) was calculated using Equation 1 for both CB₁ and CB₂, where “A” represents the compound and “CP” represents the reference compound CP55,940. The latter was appropriate to use for this selectivity determination as it is considered a non-selective cannabinoid agonist.⁵⁴

$$RA_i = \frac{E_{\max,A} \times EC_{50,CP}}{EC_{50,A} \times E_{\max,CP}} \quad (1)$$

Both RA_i values were incorporated in Equation (2), yielding a numerical “receptor selectivity factor,” calculated to assess a potential preference towards either cannabinoid receptor.

$$\text{Receptor selectivity} = \text{Log} \left(\frac{RA_i^{CB_1}}{RA_i^{CB_2}} \right) \quad (2)$$

2.4 | Analytical characterization of the OXIZID standards and the seized BZO-4en-POXIZID

The reference standards of BZO-HEXOXIZID, BZO-POXIZID, 5F-BZO-POXIZID, BZO-4en-POXIZID, and BZO-CHMOXIZID and the seized powder were analytically characterized via high-performance liquid chromatography coupled to diode-array detection (HPLC-DAD), gas chromatography coupled to mass spectrometry (GC-MS), and liquid chromatography coupled to time-of-flight mass spectrometry (LC-QTOF-MS) as described before.⁵⁵ The obtained spectra are provided in Supporting Information. For the BZO-4en-POXIZID powder, also Fourier transform infrared spectroscopy (FTIR) and nuclear magnetic resonance spectroscopy (NMR) was carried out. A short summary of each technique is provided below.

2.4.1 | High-performance liquid chromatography coupled to diode-array detection (HPLC-DAD)

Reversed-phase separation of the sample was performed on a LaChrom HPLC system from Merck-Hitachi (Tokyo, Japan), using a Merck Purospher® Star RP-8 endcapped column (5 μm , 125 mm \times 4.6 mm) with a Merck Purospher® Star RP-8 endcapped guard column (5 μm , 4 mm \times 4 mm). A diode-array detector was used to monitor a wavelength from 220 to 350 nm with a slit of 1 nm, a spectral bandwidth of 1 nm, and a spectral interval of 200 ms. The selected wavelength, used to display the chromatographic trace, was 230 nm. A total of $\sim 1 \mu\text{g}$ was injected onto the column (50 μl). For more detailed settings, the reader is referred to Supporting Information.

2.4.2 | Gas chromatography coupled to mass spectrometry (GC-MS)

One microliter of a 1-mg/ml solution was injected on an Agilent 7890A GC system coupled to a 5975 XL mass-selective detector operated by MSD Chemstation software. A 30 m \times 0.25 mm i. d. \times 0.25 μm Agilent HP-5-MS column was used. Splitless injections were performed automatically at an injection temperature of 250 $^{\circ}\text{C}$ and purge time of 1 min, with helium as a carrier gas at constant flow rate (1 ml/min). The temperature program started at 80 $^{\circ}\text{C}$ for 1 min, followed by an increase at 20 $^{\circ}\text{C}/\text{min}$ to 200 $^{\circ}\text{C}$. The temperature was then raised by 4 $^{\circ}\text{C}/\text{min}$ to 260 $^{\circ}\text{C}$ and by 30 $^{\circ}\text{C}/\text{min}$ to 300 $^{\circ}\text{C}$, which was held for an additional 8 min. Transfer line temperature and ion

source temperature were set at 300 $^{\circ}\text{C}$ and 230 $^{\circ}\text{C}$, respectively. The MS quadrupole temperature was set at 150 $^{\circ}\text{C}$, and an ionization energy of 70 eV was used. The mass spectrometer operated in SCAN mode, scanning a range of 50 to 700 m/z .

2.4.3 | Liquid chromatography coupled to time-of-flight mass spectrometry (LC-QTOF-MS)

Chromatographic separation was performed using an Agilent 1290 Infinity LC system equipped with a Phenomenex Kinetex C18-column (2.6 μm , 3 \times 50 mm), maintained at 30 $^{\circ}\text{C}$. The high-resolution mass spectrometry (HRMS) system used was a 5600 + QTOF with an electrospray ionization (ESI) source (Sciex). Upon selection of the parent compound in the quadrupole (based on mass-to-charge ratio), fragmentation occurs in the collision cell (collision energy: 35 V). Sciex Analyst TF 1.7.1 software was used to steer the system. Exact settings were the same as reported before^{55,56} and resulted in a TOF-MS full scan combined with a data dependent acquisition of product ion spectra. For more detailed settings, the reader is referred to Supporting Information.

2.4.4 | Fourier transform infrared spectroscopy (FTIR)

FTIR analysis was performed directly on the powder as received, using an Alpha-FTIR instrument from Bruker (Billerica, MA, US), equipped with an attenuated total reflection (ATR) unit. A series of 24 scans were recorded in the 400–4000 cm^{-1} wave number range, with a resolution of 4 cm^{-1} .

2.4.5 | Nuclear magnetic resonance spectroscopy (NMR)

NMR analyses were performed as previously described.⁵⁷ NMR spectra were acquired on a Bruker (Rheinstetten, Germany) spectrometer Avance II HD 600 (nominal proton frequency 600.13 MHz), equipped with a 5-mm QCI cryo-probe (^1H , ^{13}C , ^{15}N , and ^{19}F), in DMSO- d_6 solvent at 300 K. ^1H and ^{13}C NMR chemical shifts are expressed in δ scale (ppm) and referenced to the solvent (DMSO- d_6) residuals, at 2.50 and 39.52 ppm, respectively. The seized BZO-4en-POXIZID powder was characterized by one-dimensional ^1H , ^{13}C , and APT as well as $^1\text{H}/^1\text{H}$ COSY, $^1\text{H}/^1\text{H}$ TOCSY, $^1\text{H}/^{13}\text{C}$ HMBC, and $^{15}\text{N}/^1\text{H}$ HMBC experiments.

3 | RESULTS AND DISCUSSION

The activity of the novel SCRA BZO-HEXOXIZID and four structural analogs was evaluated using two similar bioassays based on the NanoBIT® technique, monitoring the recruitment of $\beta\text{arr}2$ to

either CB₁ or CB₂, upon receptor activation. This event typically results in desensitization and internalization of the receptor, thereby preventing further G protein-mediated downstream signaling. Compared to other commercialized β arr2 recruitment assays, such as the PathHunter[®] assay (Discoverx) or Tango[™] assay (ThermoFisher Scientific) which only allow for one end-point

measurement, the NanoBiT[®] assay output covers a 2-h luminescence measurement period, taking into account the complete receptor activation profile for further calculations.⁵⁸ EC₅₀ values, representing potency, and E_{max} values, representing efficacy, are depicted in Table 2. Concentration–response curves of the compounds can be found in Figure 3.

TABLE 2 Potency (EC₅₀) and efficacy (E_{max} relative to CP55,940) values and assessment of cannabinoid receptor selectivity of BZO-HEXOXIZID and analogs at either the CB₁ or CB₂ receptor

| Compound | CB ₁ | | CB ₂ | | Ratio of potencies CB ₁ /CB ₂ | Receptor selectivity ^a (CB ₁ /CB ₂) |
|-------------------------------|--------------------------------|-------------------------------|--------------------------------|-------------------------------|--|--|
| | EC ₅₀ (nM) (95% CI) | E _{max} (%) (95% CI) | EC ₅₀ (nM) (95% CI) | E _{max} (%) (95% CI) | | |
| BZO-HEXOXIZID | 721 (428–1192) | 165 (149–180) | 25.9 (10.0–67.5) | 35.0 (31.0–39.1) | 27.8 | –0.62 |
| BZO-POXIZID | 244 (142–420) | 686 (609–768) | 12.2 (3.95–40.1) | 59.8 (51.8–68.2) | 20.0 | –0.09 |
| 5F-BZO-POXIZID | 226 (136–378) | 731 (657–810) | 4.11 (0.86–18.0) | 51.7 (40.9–63.5) | 55.0 | –0.44 |
| BZO-4en-POXIZID | 532 (227–1192) | 399 (328–480) | 12.6 (2.53–63.1) | 54.1 (43.3–65.7) | 42.2 | –0.61 |
| Seized powder BZO-4en-POXIZID | 521 (300–882) | 318 (280–359) | 14.5 (2.20–97.7) | 54.1 (41.5–67.9) | 35.9 | –0.64 |
| BZO-CHMOXIZID | 84.6 (23–275) | 716 (566–876) | 2.21 (0.72–7.03) | 69.2 (59.7–79.2) | 38.3 | –0.42 |
| JWH-018 | 23.9 (11.3–52.9) | 340 (306–376) | 6.78 (2.93–14.9) | 74.0 (66.7–81.4) | 3.53 | –0.27 |
| CP55,940 | 0.69 (0.25–1.74) | 99.7 (87.5–112) | 0.49 (0.16–1.37) | 100 (87.4–113) | 1.41 | 0 |

^aReceptor selectivity calculated in a way similar to bias calculation, using relative intrinsic activities.

Note: Particular potency and efficacy values in italics belong to a seized sample (not a pure reference standard).




FIGURE 3 Activation profiles obtained for BZO-HEXOXIZID and analogs, JWH-018 and reference compound CP55,940 at the CB₁ receptor (A) and the CB₂ receptor (B). Panels (C) and (D) depict activation profiles at the CB₁ receptor (C) and CB₂ receptor (D) of the seized BZO-4en-POXIZID powder, compared with the standard of the BZO-4en-POXIZID, JWH-018, and the reference compound CP55,940. Each datapoint represents the mean \pm standard error of the mean (SEM). All data were normalized to the maximal response of CP55,940, arbitrarily set at 100% [Colour figure can be viewed at wileyonlinelibrary.com]

3.1 | Structure–activity relationship at CB₁ and CB₂

At CB₁, all tested compounds were found to be full agonists in comparison with the reference compound CP55,940, with E_{max} values exceeding 100%. On the other hand, all compounds behaved as partial agonists at CB₂, compared to the reference, with relative efficacies ranging from 35.0 to 69.2%.

In both assays, BZO-HEXOXIZID exhibited the lowest potency and efficacy of the analyzed set. The EC₅₀ and E_{max} values found in the CB₁-βarr2 assay were 721 nM and 165%, respectively, whereas at CB₂, an EC₅₀ of 25.9 nM and E_{max} of 35.0% were calculated. Shortening the n-hexyl tail to an n-pentyl tail resulted in a substantial increase in both potency and relative efficacy at CB₁, with BZO-POXIZID showing an EC₅₀ value of 244 nM and E_{max} value of 686%. Its terminally fluorinated counterpart 5F-BZO-POXIZID showed very similar activation profiles, with an EC₅₀ of 226 nM and an E_{max} of 731%. Our data are in line with those reported by Diaz et al., who, using a [³⁵S] GTP-γ-S assay, compared the CB₁ activation potential of BZO-HEXOXIZID with that of BZO-POXIZID and also observed a higher functional activity for the latter.²⁷ The absence of an impact of fluorination on intrinsic CB₁ activation potential is in line with previous findings observed for CUMYL-PEGACLONE and its 5F analog.⁴⁹ Looking at CB₂ activation, a slight increase in potency can be noticed for 5F-BZO-POXIZID (EC₅₀ = 4.11 nM) relative to BZO-POXIZID (EC₅₀ = 12.2 nM), although this difference is relatively minor. Relative efficacies at CB₂ of both analogs were also in the same order of magnitude, that is, 51.7% for 5F-BZO-POXIZID and 59.8% for BZO-POXIZID.

Furthermore, the CB₁ activation data suggest that the presence of a double bond in the pentyl tail, as present in BZO-4en-POXIZID, negatively impacted both the potency and efficacy (EC₅₀ = 532 nM, E_{max} = 399%), relative to BZO-POXIZID. Interestingly, at CB₂, this negative impact could not be demonstrated, with very similar EC₅₀ and E_{max} values for BZO-4en-POXIZID and BZO-POXIZID (12.6 vs. 12.2 nM and 54.1% vs. 59.8%, respectively). However, this decrease in activity is not unequivocally reflected in literature. When comparing JWH-018 with its unsaturated analog JWH-022, the latter was found to be more potent *in vivo*, as demonstrated via monitoring of antinociception, hypomotility, hypothermia, catalepsy in mice, and discriminative stimulus effects in rats, indicating that a pentenyl tail does not universally have a negative effect on cannabinoid activity.⁵⁹ Furthermore, using our CB₁-βarr2 assay, we previously found that the unsaturated MDMB-4en-PICA had roughly the same potency and efficacy at CB₁ (EC₅₀ = 3.70 nM, E_{max} = 289%) as its fluorinated, saturated analog 5F-MDMB-PICA (EC₅₀ = 2.13 nM, E_{max} = 289%).⁶⁰

The most potent OXIZID SCRA of this set in terms of both CB₁ and CB₂ activation was BZO-CHMOXIZID, with EC₅₀ values of 84.6 and 2.21 nM, respectively. With an E_{max} of 716% compared with the reference, its efficacy at CB₁ lies within the same range of that of BZO-POXIZID and 5F-BZO-POXIZID. The E_{max} value obtained at CB₂ was 69.2%, which ranks it among the most efficacious SCRA of this set. These findings align well with those of Diaz et al., who reported

that replacing the aliphatic tail of BZO-HEXOXIZID (MDA-19) by a cyclohexyl methyl resulted in an important increase in activity at both receptors in a [³⁵S]GTP-γ-S assay, yielding the most potent compound of the analyzed set.²⁷ Looking at other SCRA, however, this is not a consistent finding. For instance, comparing the CB₁ activity of the assumed SCRA intermediate>NNL-3 (HOBt-5F-P7AIC, carrying a fluoro pentyl tail), with its defluorinated (HOBt-P7AIC) or its cyclohexyl methyl (HOBt-CHM7AIC) analog, we noticed a dramatic decrease in activity for the HOBt-CHM7AIC, whereas both pentyl analogs had a quite similar activation profile.⁶¹ Furthermore, when comparing two L-valine SCRA at CB₁, replacement of the fluoro pentyl tail in 5F-AB-PINACA (EC₅₀ = 55.4 nM) by a cyclohexyl methyl moiety in AB-CHMINACA (EC₅₀ = 3.45 nM) did yield a more potent compound, whereas the *tert*-leucine analogs 5F-MDMB-PINACA (EC₅₀ = 0.84 nM) and MDMB-CHMINACA (EC₅₀ = 0.78 nM) had essentially the same potency at CB₁.⁵⁰

It is interesting to highlight that, although we found these compounds to exhibit a broad range of intrinsic activities at CB₁, Diaz et al. did not observe large differences in binding affinity of BZO-HEXOXIZID, BZO-POXIZID, and BZO-CHMOXIZID for both cannabinoid receptors, as evaluated using radioligand binding assays.²⁷ This emphasizes the fact that the differences in activities, as demonstrated in our bioassays, are most likely not the consequence of different receptor affinities, but rather the result of other or better interactions with residues inside the binding pocket of the receptors. Compared with the efficacies obtained using the CB₁-βarr2 bioassay, E_{max} values for the CB₂ receptor are less divergent, which is in agreement with past analyses in which we have consistently noticed a more clustered profile for CB₂. To this day, the underlying reason for this has not been elucidated.

In summary, a general trend could be noticed regarding the impact of the tail of these OXIZID SCRA on potency and relative efficacy at CB₁. BZO-HEXOXIZID, carrying a hexyl tail, had the lowest activity, followed by the 4-pentenyl analog BZO-4en-POXIZID. Although the saturated BZO-POXIZID and its fluoro pentyl analog 5F-BZO-POXIZID were more potent and efficacious, the lowest EC₅₀ value (and hence highest potency) was observed for BZO-CHMOXIZID, the only SCRA in this set carrying a cyclic tail. Overall, at CB₂, the same rank order in terms of potency was applicable, albeit less distinct.

3.2 | Assessment of cannabinoid receptor selectivity

BZO-HEXOXIZID was originally selected as a potential lead compound in the search for new therapeutics for neuropathic pain, based on its potency at CB₂ (63.4 nM), its CB₂ selectivity and its only moderate potency at CB₁, as assessed by means of a [³⁵S] guanosine-5'-triphosphate (GTP)-γ-S assay.²⁷ To further investigate the receptor preference (CB₁ vs. CB₂) of BZO-HEXOXIZID and its analogs, two methods were implemented. First, in line with the method applied by Banister et al., the ratio of EC₅₀ values (CB₁/CB₂) was calculated.⁶² A

higher value reflects a larger fold difference between both EC_{50} values, indicative of a more CB_2 selective compound. Second, and more elaborately, a numerical value, similar to a bias factor, was calculated. This calculation entails the relative intrinsic activity (RA_i), which takes into account both the EC_{50} (potency) and E_{max} (efficacy) value of a compound at the two cannabinoid receptors. As this method includes both pharmacological parameters (potency and efficacy) in the equation and therefore considers multiple aspects of CB_1 and CB_2 activation, it may be a more comprehensive and complete approach to evaluate receptor selectivity. Values below 0 indicate a preference towards CB_2 and therefore a potential CB_2 selectivity.

Implementing the EC_{50} ratio method, all SCRA exhibited a clear CB_2 selectivity (20.0–55.0), compared with CP55,940 (1.41) and JWH-018 (3.53). The same conclusion could be drawn from the bias formula method, assigning a preference towards CB_2 activation for all compounds, except for the n-pentyl analog BZO-POXIZID. In fact, BZO-POXIZID was found to be the least CB_2 selective using both calculation methods. Taken together, the CB activation profile of most OXIZID compounds somewhat resembles that of XLR-11, a SCRA found in authentic urine samples of drug users in 2017, which also demonstrated a CB_2 preference in our bioassay.⁴⁶ Overall, the assessment of receptor selectivity based on the formula also implemented for bias calculation seems to present a clearer view on the selective behavior of these substances, as differences appear to be more pronounced compared with the somewhat clustered EC_{50} ratios.

3.3 | Analytical characterization of a seized powder containing BZO-4en-POXIZID

A yellow powder, in which the presence of BZO-4en-POXIZID was demonstrated, was intercepted by the Belgian Customs in November 2021. This powder was characterized alongside the reference standard for BZO-4en-POXIZID using HPLC-DAD, GC-MS, and LC-QTOF-MS, as well as with FTIR and NMR, which confirmed the identity of the powder (Supporting Information), with no impurities being detected. This powder was also analyzed alongside the BZO-4en-POXIZID reference standard for its activity in the CB_1 and CB_2 bioassays. Figure 3C,D illustrates the similar activation profiles for the BZO-4en-POXIZID powder and the reference standard. Given the comparable potency and efficacy at both receptors (EC_{50} 512 nM, E_{max} 318% for the powder vs. EC_{50} 532 nM, E_{max} 399% for the BZO-4en-POXIZID standard at CB_1), a high level of purity of the powder can be assumed, in line with the analytical characterization.

3.4 | Analytical characterization of reference standards of a panel of OXIZID SCRA

Similar to the analysis of the seized sample, reference standards of BZO-HEXOXIZID, BZO-POXIZID, 5F-BZO-POXIZID, BZO-4en-POXIZID, and BZO-CHMOXIZID were characterized using HPLC-DAD, GC-MS, and LC-QTOF-MS. Results were in line with findings

reported by Liu et al.,²⁰ who characterized BZO-POXIZID and 5F-BZO-POXIZID using GC-MS and QTOF-MS. For BZO-HEXOXIZID and BZO-CHMOXIZID, results were in accordance with the analytical reports distributed by the CFSRE.^{36,39} Chromatograms and mass spectra can be found in Supporting Information.

4 | CONCLUSION

This study is the first to report on the in vitro intrinsic receptor activation potential at CB_1 and CB_2 of the newly emerging SCRA BZO-HEXOXIZID and four structural analogs. Using two live cell-based β arr2 recruitment assays, all compounds were found to be full agonists at CB_1 , with efficacies ranging from 165% to 731% compared with CP55,940, and with potencies (EC_{50}) ranging from 84.6 (BZO-CHMOXIZID) to 721 nM (BZO-HEXOXIZID), all being less potent than CP55,940. The n-hexyl analog BZO-HEXOXIZID (also known in literature as MDA-19) had the lowest potency and efficacy, followed by the pentenyl analog BZO-4en-POXIZID. Shortening the n-hexyl tail resulted in an important increase in CB_1 activation potential. The pentyl and fluoro pentyl analogs BZO-POXIZID and 5F-BZO-POXIZID exhibited higher but quite similar potencies, demonstrating that the addition of a fluorine atom did not have a major impact on CB_1 activation. The most potent SCRA of the investigated set was the cyclohexyl methyl analog BZO-CHMOXIZID, which had a relative efficacy within the same range as that of BZO-POXIZID and 5F-BZO-POXIZID. Overall, the same general trend and rank order regarding potency was seen at CB_2 , although differences were less pronounced. More specifically, the negative impact of an unsaturated hydrocarbon tail was not observed at CB_2 . All OXIZIDs showed a clear preference for CB_2 , compared with CP55,940. Given the rather moderate potencies found for these compounds at CB_1 , it is premature to predict whether they will pose extensive cannabinoid-related toxicity. However, these findings may be of value for drug policy makers and health care workers, as they give an idea on the pharmacology of these newly emerging SCRA and may hint at substances that could potentially appear in the future. Depending on multiple variables such as ease of synthesis, price and availability, it still remains to be seen whether and to what extent this new class will “take off” on the recreational drug market.

ACKNOWLEDGEMENTS

M.H.D acknowledges funding from the Research Foundation-Flanders (FWO; Grant 1S54521N). Cayman Chemical is acknowledged for providing the tested compounds. I.M.J.V acknowledges analytical data sharing by the European Commission Directorate General for Taxation and Customs Unions (DG TAXUD) and the Joint Research Centre (JRC) (Administrative Arrangement JRC-Nr35320-CLEN2SAND3-DG TAXUD/2018/DE/334 for fast recognition of New Psychoactive Substances (NPS) and identification of unknown chemicals). We gratefully acknowledge the lab technicians of the Laboratory of Toxicology for performing part of the analytical characterizations. Margaret Holland is acknowledged for the technical assistance for the NMR analysis on the seized BZO-4en-POXIZID sample.


DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID

Marie H. Deventer  <https://orcid.org/0000-0001-6667-2561>

Katleen Van Uytvanghe  <https://orcid.org/0000-0001-8195-150X>

Inge M. J. Vinckier  <https://orcid.org/0000-0003-2394-3485>

Fabiano Reniero  <https://orcid.org/0000-0002-0915-8168>

Claude Guillou  <https://orcid.org/0000-0002-6210-3457>

Christophe P. Stove  <https://orcid.org/0000-0001-7126-348X>

REFERENCES

- United Nations Office on Drugs and Crime. World drug report 2021. 2021. Accessed November 24, 2021. <https://www.unodc.org/unodc/en/data-and-analysis/wdr2021.html>
- European Monitoring Centre for Drugs and Drug Addiction, ed. *European Drug Report: Trends and Developments 2020*. Publications Office of the European Union; 2020. https://www.emcdda.europa.eu/system/files/publications/13236/TDAT20001ENN_web.pdf
- Duke K. Producing the 'problem' of new psychoactive substances (NPS) in English prisons. *Int J Drug Policy*. 2020;80. doi:10.1016/j.drugpo.2019.05.022
- Gray P, Ralphs R, Williams L. The use of synthetic cannabinoid receptor agonists (SCRAs) within the homeless population: motivations, harms and the implications for developing an appropriate response. *Addict Res Theory*. 2021;29(1):1-10. doi:10.1080/16066359.2020.1730820
- Seely KA, Lapoint J, Moran JH, Fattore L. Spice drugs are more than harmless herbal blends: a review of the pharmacology and toxicology of synthetic cannabinoids. *Prog Neuropsychopharmacol Biol Psychiatry*. 2012;39(2):234-243. doi:10.1016/j.pnpbp.2012.04.017
- Lapoint J, James LP, Moran CL, Nelson LS, Hoffman RS, Moran JH. Severe toxicity following synthetic cannabinoid ingestion. *Clin Toxicol Phila Pa*. 2011;49(8):760-764. doi:10.3109/15563650.2011.609822
- Kemp AM, Clark MS, Dobbs T, Galli R, Sherman J, Cox R. Top 10 facts you need to know about synthetic cannabinoids: not so nice spice. *Am J Med*. 2016;129(3):240, e1-244. doi:10.1016/j.amjmed.2015.10.008
- European Monitoring Centre for Drugs and Drug Addiction. Perspectives on drugs: synthetic cannabinoids in Europe. 2017. Accessed March 21, 2022. https://www.emcdda.europa.eu/system/files/publications/2753/POD_Synthetic%20cannabinoids_0.pdf
- Labay LM, Caruso JL, Gilson TP, et al. Synthetic cannabinoid drug use as a cause or contributory cause of death. *Forensic Sci Int*. 2016;260:31-39. doi:10.1016/j.forsciint.2015.12.046
- Manini AF, Krotulski AJ, Schimmel J, et al. Respiratory failure in confirmed synthetic cannabinoid overdose. *Clin Toxicol*. 2021;60(4):524-526. doi:10.1080/15563650.2021.1975734
- Bukke VN, Archana M, Villani R, Serviddio G, Cassano T. Pharmacological and toxicological effects of phytocannabinoids and recreational synthetic cannabinoids: increasing risk of public health. *Pharmaceuticals*. 2021;14(10):965. doi:10.3390/ph14100965
- Riederer AM, Campleman SL, Carlson RG, et al. Acute poisonings from synthetic cannabinoids—50 U.S. toxicology investigators consortium registry sites, 2010–2015. *MMWR Morb Mortal Wkly Rep*. 2016;65(27):692-695. doi:10.15585/mmwr.mm6527a2
- Abouchedid R, Hudson S, Thurtle N, et al. Analytical confirmation of synthetic cannabinoids in a cohort of 179 presentations with acute recreational drug toxicity to an Emergency Department in London, UK in the first half of 2015. *Clin Toxicol*. 2017;55(5):338-345. doi:10.1080/15563650.2017.1287373
- Office of China National Narcotics Control Commission. Announcement on the inclusion of 18 substances including synthetic cannabinoids and flumaine in the Supplementary List of Controlled Narcotic Drugs and Psychotropic Substances with Non-medical Use. Published online May 12, 2021. <https://app.mps.gov.cn/gdnps/pc/content.jsp?id=7881703>
- The Center for Forensic Science Research & Education, Cayman Chemical. New systematic naming for synthetic cannabinoid "MDA-19" and its related analogues: BZO-HEXOXIZID, 5F-BZO-POXIZID, and BZO-POXIZID. Published online August 31, 2021. Accessed November 25, 2021. https://www.npsdiscovery.org/wp-content/uploads/2021/08/New-Systematic-Naming-for-MDA-19-and-Related-Analogues_NPS-Discovery_083121.pdf
- Cayman Chemical. BZO-4en-POXIZID product information. Published online November 11, 2021. Accessed December 1, 2021. <https://cdn.caymanchem.com/cdn/insert/35503.pdf>
- van Amsterdam J, Nutt D, van den Brink W. Generic legislation of new psychoactive drugs. *J Psychopharmacol (Oxf)*. 2013;27(3):317-324. doi:10.1177/0269881112474525
- United Nations Office on Drugs and Crime. News: April 2019—China: announcement to place all fentanyl-related substances under national control. Published April 2019. Accessed November 25, 2021. <https://www.unodc.org/LSS/Announcement/Details/f2adea68-fbed-4292-a4cc-63771c943318>
- United Nations Office on Drugs and Crime. News: May 2021—China: announcement to place synthetic cannabinoids under generic control. Published May 2021. Accessed November 24, 2021. <https://www.unodc.org/LSS/Announcement/Details/ff032a29-2e14-4dab-b7d8-ab86d355c809>
- Liu CM, Hua ZD, Jia W, Li T. Identification of AD-18, 5F-MDA-19, and pentyl MDA-19 in seized materials after the class-wide ban of synthetic cannabinoids in China. *Drug Test Anal*. Published online October 25, 2021. doi:10.1002/dta.3185
- Li X, Shen L, Hua T, Liu ZJ. Structural and functional insights into cannabinoid receptors. *Trends Pharmacol Sci*. 2020;41(9):665-677. doi:10.1016/j.tips.2020.06.010
- Turcotte C, Blanchet MR, Laviolette M, Flamand N. The CB2 receptor and its role as a regulator of inflammation. *Cell Mol Life Sci*. 2016;73(23):4449-4470. doi:10.1007/s00018-016-2300-4
- Guindon J, Hohmann AG. Cannabinoid CB2 receptors: a therapeutic target for the treatment of inflammatory and neuropathic pain. *Br J Pharmacol*. 2008;153(2):319-334. doi:10.1038/sj.bjp.0707531
- Hashiesh HM, Sharma C, Goyal SN, Jha NK, Ojha S. Pharmacological properties, therapeutic potential and molecular mechanisms of JWH133, a CB2 receptor-selective agonist. *Front Pharmacol*. 2021;12:1818. doi:10.3389/fphar.2021.702675
- Bie B, Wu J, Foss JF, Naguib M. An overview of the cannabinoid type 2 (CB2) receptor system and its therapeutic potential. *Curr Opin Anaesthesiol*. 2018;31(4):407-414. doi:10.1097/ACO.0000000000000616
- Malan TP, Ibrahim MM, Lai J, Vanderah TW, Makriyannis A, Porreca F. CB2 cannabinoid receptor agonists: pain relief without psychoactive effects? *Curr Opin Pharmacol*. 2003;3(1):62-67. doi:10.1016/S1471-4892(02)00004-8
- Diaz P, Xu J, Astruc-Diaz F, Pan HM, Brown DL, Naguib M. Design and synthesis of a novel series of N-alkyl isatin acylhydrazones derivatives that act as selective cannabinoid receptor 2 agonists for the treatment of neuropathic pain. *J Med Chem*. 2008;51(16):4932-4947. doi:10.1021/jm8002203
- Xu JJ, Diaz P, Astruc-Diaz F, Craig S, Munoz E, Naguib M. Pharmacological characterization of a novel cannabinoid ligand, MDA19, for treatment of neuropathic pain. *Anesth Analg*. 2010;111(1):99-109. doi:10.1213/ANE.0b013e3181e0cdaf

29. St. John Smith E. Advances in understanding nociception and neuro-pathic pain. *J Neurol.* 2018;265(2):231-238. doi:10.1007/s00415-017-8641-6
30. Jensen TS, Gottrup H, Sindrup SH, Bach FW. The clinical picture of neuropathic pain. *Eur J Pharmacol.* 2001;429(1):1-11. doi:10.1016/S0014-2999(01)01302-4
31. Baron R, Binder A, Wasner G. Neuropathic pain: diagnosis, patho-physiological mechanisms, and treatment. *Lancet Neurol.* 2010;9(8):807-819. doi:10.1016/S1474-4422(10)70143-5
32. Rao M, Chen D, Zhan P, Jiang J. MDA19, a novel CB2 agonist, inhibits hepatocellular carcinoma partly through inactivation of AKT signaling pathway. *Biol Direct.* 2019;14(1):9. doi:10.1186/s13062-019-0241-1
33. Liu B, Xu L, Dai EN, Tian JX, Li JM. Anti-tumoral potential of MDA19 in human osteosarcoma via suppressing PI3K/Akt/mTOR signaling pathway. *Biosci Rep.* 2018;38(6):BSR20181501. doi:10.1042/BSR20181501
34. Dang N, Meng X, Ma S, et al. MDA-19 suppresses progression of melanoma via inhibiting the PI3K/Akt pathway. *Open Med.* 2018;13(1):416-424. doi:10.1515/med-2018-0061
35. European Monitoring Centre for Drugs and Drug Addiction. EMCDDA-Europol 2016 Annual Report on the Implementation of Council Decision 2005/387/JHA. 2016. https://www.emcdda.europa.eu/system/files/publications/4724/TDAN17001ENN_PDFWEB.pdf
36. The Center for Forensic Science Research & Education. BZO-HEXOXIZID chemistry report. 2021. Accessed November 25, 2021. https://www.npsdiscovery.org/wp-content/uploads/2021/10/BZO-HEXOXIZID_101921_CFSRE-Chemistry_Report.pdf?mc_cid=c736522745&mc_eid=da2366d692
37. The Center for Forensic Science Research & Education. 5F-BZO-POXIZID chemistry report. 2021. Accessed November 25, 2021. https://www.npsdiscovery.org/wp-content/uploads/2021/10/5F-BZO-POXIZID_101921_CFSRE-Chemistry_Report.pdf?mc_cid=c736522745&mc_eid=da2366d692
38. The Center for Forensic Science Research & Education. BZO-POXIZID chemistry report. 2021. Accessed November 25, 2021. https://www.npsdiscovery.org/wp-content/uploads/2021/10/BZO-POXIZID_101921_CFSRE-Chemistry_Report.pdf?mc_cid=c736522745&mc_eid=da2366d692
39. The Center for Forensic Science Research & Education. BZO-CHMOXIZID chemistry report. 2021. Accessed November 25, 2021. https://www.npsdiscovery.org/wp-content/uploads/2021/11/BZO-CHMOXIZID_111821_CFSRE-Chemistry_Report.pdf?mc_cid=0ee18051f6&mc_eid=da2366d692
40. The Center for Forensic Science Research & Education. 2021 Q1 Synthetic cannabinoids trend report. 2021. Accessed December 1, 2021. https://www.npsdiscovery.org/wp-content/uploads/2021/04/2021-Q1_Synthetic-Cannabinoids_Trend-Report.pdf
41. The Center for Forensic Science Research & Education. 2021 Q2 Synthetic cannabinoids trend report. 2021. Accessed December 1, 2021. https://www.npsdiscovery.org/wp-content/uploads/2021/07/2021-Q2_Synthetic-Cannabinoids_Trend-Report.pdf
42. The Center for Forensic Science Research & Education. 2021 Q3 Synthetic cannabinoids trend report. 2021. Accessed December 1, 2021. https://www.npsdiscovery.org/wp-content/uploads/2021/10/2021-Q3_Synthetic-Cannabinoids_Trend-Report.pdf
43. Ministry of Justice. Notification 2021/0903/HU: Amendment of Decree No 55/2014 of the Ministry of Human Capacities of 30 December 2014 on substances or groups of compounds classified as new psychoactive substances. 2021. Accessed January 5, 2022. <https://ec.europa.eu/growth/tools-databases/tris/en/search/?trisaction=search.detail&year=2021&num=903>
44. Ministry of Justice. Notification 2021/0840/HU: Amendment of Decree No 55/2014 of the Ministry of Human Capacities of 30 December 2014 on substances or groups of compounds classified as new psychoactive substances. 2021. Accessed January 10, 2022. <https://ec.europa.eu/growth/tools-databases/tris/index.cfm/en/search/?trisaction=search.detail&year=2021&num=840&mLang=EN>
45. Cannaert A, Storme J, Franz F, Auwärter V, Stove CP. Detection and activity profiling of synthetic cannabinoids and their metabolites with a newly developed bioassay. *Anal Chem.* 2016;88(23):11476-11485. doi:10.1021/acs.analchem.6b02600
46. Cannaert A, Franz F, Auwärter V, Stove CP. Activity-based detection of consumption of synthetic cannabinoids in authentic urine samples using a stable cannabinoid reporter system. *Anal Chem.* 2017;89(17):9527-9536. doi:10.1021/acs.analchem.7b02552
47. Cannaert A, Storme J, Hess C, Auwärter V, Wille SMR, Stove CP. Activity-based detection of cannabinoids in serum and plasma samples. *Clin Chem.* 2018;64(6):918-926. doi:10.1373/clinchem.2017.285361
48. Cannaert A, Sparkes E, Pike E, et al. Synthesis and in vitro cannabinoid receptor 1 activity of recently detected synthetic cannabinoids 4F-MDMB-BICA, 5F-MPP-PICA, MMB-4en-PICA, CUMYL-CBMICA, ADB-BINACA, APP-BINACA, 4F-MDMB-BINACA, MDMB-4en-PINACA, A-CHMINACA, 5F-AB-P7AICA, 5F-MDMB-P7AICA, and 5F-AP7AICA. *ACS Chem Neurosci.* 2020;11(24):4434-4446. doi:10.1021/acscchemneuro.0c00644
49. Janssens L, Cannaert A, Connolly MJ, Liu H, Stove CP. In vitro activity profiling of Cumyl-PEGACLONE variants at the CB1 receptor: fluorination versus isomer exploration. *Drug Test Anal.* 2020;12(9):1336-1343. doi:10.1002/dta.2870
50. Wouters E, Mogler L, Cannaert A, Auwärter V, Stove C. Functional evaluation of carboxy metabolites of synthetic cannabinoid receptor agonists featuring scaffolds based on L-valine or L-tert-leucine. *Drug Test Anal.* 2019;11(8):1183-1191. doi:10.1002/dta.2607
51. Rajagopal S, Ahn S, Rominger DH, et al. Quantifying ligand bias at seven-transmembrane receptors. *Mol Pharmacol.* 2011;80(3):367-377. doi:10.1124/mol.111.072801
52. Pottie E, Dedecker P, Stove CP. Identification of psychedelic new psychoactive substances (NPS) showing biased agonism at the 5-HT2AR through simultaneous use of β -arrestin 2 and miniG α q bioassays. *Biochem Pharmacol.* 2020;182:114251. doi:10.1016/j.bcp.2020.114251
53. Vandeputte MM, Van Uytfaenge K, Layle NK, St. Germaine DM, Iula DM, Stove CP. Synthesis, chemical characterization, and μ -opioid receptor activity assessment of the emerging group of "nitazene" 2-benzylbenzimidazole synthetic opioids. *ACS Chem Neurosci.* 2021;12(7):1241-1251. doi:10.1021/acscchemneuro.1c00064
54. Felder CC, Joyce KE, Briley EM, et al. Comparison of the pharmacology and signal transduction of the human cannabinoid CB1 and CB2 receptors. *Mol Pharmacol.* 1995;48(3):443-450.
55. Blanckaert P, Cannaert A, Van Uytfaenge K, et al. Report on a novel emerging class of highly potent benzimidazole NPS opioids: chemical and in vitro functional characterization of isotonitazene. *Drug Test Anal.* 2020;12(4):422-430. doi:10.1002/dta.2738
56. Thoren KL, Colby JM, Shugarts SB, Wu AHB, Lynch KL. Comparison of information-dependent acquisition on a tandem quadrupole TOF vs a triple quadrupole linear ion trap mass spectrometer for broad-spectrum drug screening. *Clin Chem.* 2016;62(1):170-178. doi:10.1373/clinchem.2015.241315
57. Lobo Vicente J, Chassaing H, Holland MV, et al. Systematic analytical characterization of new psychoactive substances: a case study. *Forensic Sci Int.* 2016;265:107-115. doi:10.1016/j.forsciint.2016.01.024
58. Wouters E, Walraed J, Banister SD, Stove CP. Insights into biased signaling at cannabinoid receptors: synthetic cannabinoid receptor

- agonists. *Biochem Pharmacol.* 2019;169:113623. doi:[10.1016/j.bcp.2019.08.025](https://doi.org/10.1016/j.bcp.2019.08.025)
59. Wiley JL, Compton DR, Dai D, et al. Structure-activity relationships of indole- and pyrrole-derived cannabinoids. *J Pharmacol Exp Ther.* 1998;285(3):995-1004.
60. Antonides LH, Cannaeert A, Norman C, et al. Shape matters: the application of activity-based in vitro bioassays and chiral profiling to the pharmacological evaluation of synthetic cannabinoid receptor agonists in drug-infused papers seized in prisons. *Drug Test Anal.* 2021;13(3):628-643. doi:[10.1002/dta.2965](https://doi.org/10.1002/dta.2965)
61. Ametovski A, Cairns EA, Grafinger KE, et al. NNL-3: a synthetic intermediate or a new class of hydroxybenzotriazole esters with cannabinoid receptor activity? *ACS Chem Neurosci.* 2021;12(21):4020-4036. doi:[10.1021/acscemneuro.1c00348](https://doi.org/10.1021/acscemneuro.1c00348)
62. Banister SD, Moir M, Stuart J, et al. Pharmacology of indole and indazole synthetic cannabinoid designer drugs AB-FUBINACA, ADB-FUBINACA, AB-PINACA, ADB-PINACA, 5F-AB-PINACA, 5F-ADB-PINACA, ADBICA, and 5F-ADBICA. *ACS Chem Neurosci.* 2015;6(9):1546-1559. doi:[10.1021/acscemneuro.5b00112](https://doi.org/10.1021/acscemneuro.5b00112)

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Deventer MH, Van Uytvanghe K, Vinckier IMJ, Reniero F, Guillou C, Stove CP. Cannabinoid receptor activation potential of the next generation, generic ban evading OXIZID synthetic cannabinoid receptor agonists. *Drug Test Anal.* 2022;1-11. doi:[10.1002/dta.3283](https://doi.org/10.1002/dta.3283)